

Electromagnetic interaction of a magnetized rotating star with a conducting disk

Ya. N. Istomin¹

© Springer-Verlag ●●●●

Abstract A conducting disk significantly changes the generation of the electromagnetic radiation excited by the rotation of the magnetic field frozen to a star. Due to the reflection of waves from a disk there appear waves propagating toward a star, not only outward a star as it takes place for the magneto-dipole radiation. Because that the angular momentum can be transformed from a disk to a star when the inner edge of a disk approaches the light surface of a rotating star. This is purely electromagnetic effect. At some distance of a disk from a star, $r_d = r^* \simeq c/\omega_s$, the stellar angular momentum losses due to the electromagnetic radiation become zero. It results the stable stellar rotation.

Keywords stars: neutron

1 Introduction

The interaction of an accretion disk with a magnetized star is very complicated and is not understandable yet. The observations of X-ray binaries demonstrate many physical phenomenon [1]. It seems useful to solve a set of simple physical problems throwing light on the nature of interaction of an accretion flow with a star. One of these problems is the interaction of a rotating magnetic field with a conducting matter of a disk. The phenomena depends on the conductivity of the disk matter σ . The physics is different for the high and low conductivities. Aly [2] consider the disk conductivity is high and the magnetic field does not penetrate to a disk. Opposite, Bardou and Heyvaerts [3] consider the conductivity is low due to abnormal resistivity of a turbulent disk plasma, and the field penetrate to a disk freely.

The interesting combination of penetration and repulsion of magnetic field lines was suggested by Lovelace et. al [4]. We consider here a disk to be stationary, without accretion and with the classical properties.

The ionized plasma of an accretion disk has the conductivity $\sigma = 10^{13}(T_e/1\text{eV})^{3/2}(\Lambda/10)^{-1}\text{s}^{-1}$, which is high enough to consider a disk as an ideal conductor. Here T_e is the temperature of electrons in a disk which is larger than 10eV, Λ is the Coulomb logarithm. At such conductivity σ the width of the skin layer $\lambda_{sk} = (\tau c^2/\sigma)^{1/2}$ is less than the disk width H . The value of τ is the characteristic time of turbulent motion in the α -disk, $\tau = H/v_k$, v_k is the Keplerian velocity of the disk rotation. The condition $\sigma \gg c^2/Hv_k$ is well fulfilled in the inner parts of a disk.

Here we consider the oblique magnetic field rotating with the frequency of the star rotation ω_s , and a disk having the infinite conductivity and finite dimensions. The environment of this system is the vacuum. We will see that these conditions though the boundary condition on a disk surface results to the repulsion of the magnetic field from a disk.

2 Statement of problem

We are solving the following problem: a rotating neutron star with the angular velocity ω_s and the magnetic momentum μ is surrounded by a disk (see figure 1). The angle between ω_s and μ is χ . The matter of a disk is ideal, i.e. the conductivity $\sigma = \infty$. It means $\mathbf{E}' = 0$, where $\mathbf{E}'$ is the electric field in the frame of a disk. The field in the laboratory frame is

$$\mathbf{E} = -\frac{1}{c}[\mathbf{v}_d\mathbf{B}]. \quad (1)$$

Here $\mathbf{v}_d$ is the rotational velocity of the disk matter around the a star $\mathbf{v}_d = v_k\mathbf{e}_\phi$. We consider a disk is

Ya. N. Istomin

¹P. N. Lebedev Physical Institute, Leninsky Prospect 53, Moscow, 119991 Russia, istomin@lpi.ru

rotating with the Keplerian velocity $v_k \propto r^{-1/2}$ in the ϕ direction

$$\mathbf{v}_d = \left(\frac{GM_s}{r} \right)^{1/2} \sin \vartheta \mathbf{e}_\phi. \quad (2)$$

Here M_s is the stellar mass, G is the gravitational constant and r, ϑ, ϕ are the spherical coordinates. To create the electric field (1) the matter of a disk must polarize and produces the charge density

$$\varrho = \frac{1}{4\pi} \nabla \mathbf{E} = -\frac{v_k}{4\pi cr} \left[\frac{1}{\sin \vartheta} \frac{\partial}{\partial \vartheta} (\sin^2 \vartheta B_r) - \frac{1}{r^{1/2}} \frac{\partial}{\partial r} (r^{3/2} B_\vartheta) \sin \vartheta \right]. \quad (3)$$

The rotating charge density produces the toroidal electric current j_ϕ , $j_\phi = v_k \varrho \sin \vartheta$,

$$j_\phi = -\frac{v_k^2}{4\pi cr} \left[\frac{\partial}{\partial \vartheta} (\sin^2 \vartheta B_r) - \frac{1}{r^{1/2}} \frac{\partial}{\partial r} (r^{3/2} B_\vartheta) \sin^2 \vartheta \right].$$

We consider that a disk occupies the region $r_d < r < \infty, \pi/2 - \delta < \vartheta < \pi/2 + \delta$, where r_d is inner edge of a disk and 2δ is its width in the ϑ direction. Integrating the Maxwell equation $(\text{curl} \mathbf{B})_\phi = (4\pi/c)(j_\phi + j'_\phi)$ ($E_\phi = 0$ inside a disk) over the angle ϑ from $\pi/2 - \delta$ to $\pi/2 + \delta$ we obtain the condition connecting two values of B_r on both sides of a disk

$$B_r(\frac{\pi}{2} - \delta) - B_r(\frac{\pi}{2} + \delta) = \left(1 - \frac{v_k^2 \cos^2 \delta}{c^2} \right)^{-1} \left[-\frac{v_k^2}{c^2 r^{1/2}} \frac{\partial}{\partial r} \left(r^{3/2} \int_{\pi/2 - \delta}^{\pi/2 + \delta} \sin^2 \vartheta B_\vartheta d\vartheta \right) + \frac{\partial}{\partial r} \left(r \int_{\pi/2 - \delta}^{\pi/2 + \delta} B_\vartheta d\vartheta \right) \right] - \frac{4\pi r}{c} \int_{\pi/2 - \delta}^{\pi/2 + \delta} j'_\phi d\vartheta. \quad (4)$$

Here the current j'_ϕ is the toroidal electric current exciting in a disk by the infinitely small electric field E'_ϕ . The boundary condition (4) is the condition for the current j'_ϕ arising inside a disk. Another two boundary conditions follow from (1)

$$E_\phi(\frac{\pi}{2} - \delta) = E_\phi(\frac{\pi}{2} + \delta) = 0, \quad r \geq r_d, \quad (5)$$




Fig. 1 A rotating neutron star with a dipole magnetic field is surrounded by a disk.

$$(E_r - \frac{v_k}{c} \sin \vartheta B_\vartheta)|_{\vartheta=\pi/2-\delta} = 0, \quad r \geq r_d; \quad (6)$$

$$(E_r - \frac{v_k}{c} \sin \vartheta B_\vartheta)|_{\vartheta=\pi/2+\delta} = 0, \quad r \geq r_d.$$

3 Solution

The general solution of the wave equation in vacuum in the spherical coordinates consists of the sum of the multipole components, magnetic and electric. The magnetic multipole of the n -th order is

$$\mathbf{B}_n = \begin{cases} P_n^1(\cos \vartheta) z^{-\frac{3}{2}} [a_n^B J_{n+\frac{1}{2}} + b_n^B J_{-n-\frac{1}{2}}], & \mathbf{e}_r \\ \frac{1}{n(n+1)} \frac{\partial P_n^1}{\partial \vartheta} \frac{1}{z} \frac{\partial}{\partial z} z^{\frac{1}{2}} [a_n^B J_{n+\frac{1}{2}} + b_n^B J_{-n-\frac{1}{2}}], & \mathbf{e}_\vartheta \\ \frac{i}{n(n+1)} \frac{P_n^1}{\sin \vartheta} \frac{1}{z} \frac{\partial}{\partial z} z^{\frac{1}{2}} [a_n^B J_{n+\frac{1}{2}} + b_n^B J_{-n-\frac{1}{2}}], & \mathbf{e}_\phi, \end{cases} \quad (7)$$

$$\mathbf{E}_n = \begin{cases} 0, & \mathbf{e}_r \\ -\frac{1}{n(n+1)} \frac{P_n^1}{\sin \vartheta} z^{-\frac{1}{2}} [a_n^B J_{n+\frac{1}{2}} + b_n^B J_{-n-\frac{1}{2}}], & \mathbf{e}_\vartheta \\ -\frac{i}{n(n+1)} \frac{\partial P_n^1}{\partial \vartheta} z^{-\frac{1}{2}} [a_n^B J_{n+\frac{1}{2}} + b_n^B J_{-n-\frac{1}{2}}], & \mathbf{e}_\phi. \end{cases}$$

Here and below $P_n^1(\cos \vartheta)$ are the associated Legendre functions, $J_{n+1/2}(z)$ is the Bessel functions of the $(n+1/2)$ order, $z = r\omega_s/c$. The electric multipole of the n -th order is

$$\mathbf{E}_n = \begin{cases} P_n^1(\cos \vartheta) z^{-\frac{3}{2}} [a_n^E J_{n+\frac{1}{2}} + b_n^E J_{-n-\frac{1}{2}}], & \mathbf{e}_r \\ \frac{1}{n(n+1)} \frac{\partial P_n^1}{\partial \vartheta} \frac{1}{z} \frac{\partial}{\partial z} z^{\frac{1}{2}} [a_n^E J_{n+\frac{1}{2}} + b_n^E J_{-n-\frac{1}{2}}], & \mathbf{e}_\vartheta \\ \frac{i}{n(n+1)} \frac{P_n^1}{\sin \vartheta} \frac{1}{z} \frac{\partial}{\partial z} z^{\frac{1}{2}} [a_n^E J_{n+\frac{1}{2}} + b_n^E J_{-n-\frac{1}{2}}], & \mathbf{e}_\phi, \end{cases} \quad (8)$$

$$\mathbf{B}_n = \begin{cases} 0, & \mathbf{e}_r \\ -\frac{1}{n(n+1)} \frac{P_n^1}{\sin \vartheta} z^{-\frac{1}{2}} [a_n^E J_{n+\frac{1}{2}} + b_n^E J_{-n-\frac{1}{2}}], & \mathbf{e}_\vartheta \\ -\frac{i}{n(n+1)} \frac{\partial P_n^1}{\partial \vartheta} z^{-\frac{1}{2}} [a_n^E J_{n+\frac{1}{2}} + b_n^E J_{-n-\frac{1}{2}}], & \mathbf{e}_\phi. \end{cases}$$

All quantities $\mathbf{B}_n$ and $\mathbf{E}_n$ are oscillating functions with the frequency of the stellar rotation ω_s , $(\mathbf{B}, \mathbf{E})_n \propto e^{i\phi - i\omega_s t}$. Multipoles (7,8) contain waves travelling in the positive direction from a neutron star to the infinity, in this case $b_n = i(-1)^{n+1}a_n$ and the combination $a_n J_{n+1/2} + b_n J_{-n-1/2} = a_n H_{n+1/2}^{(1)}$ is the Hankel function of the first kind

$$H_{n+1/2}^{(1)} = e^{iz} \left(\frac{1}{2\pi z} \right)^{1/2} Z_n \left(\frac{1}{z} \right).$$

Here $Z_n(1/z)$ is the polynomial of the n -th order of the argument z^{-1} . Also for the wave propagating in the negative direction from the infinity $b_n = i(-1)^n a_n$, and we have the Hankel function of the second kind, $H_{n+1/2}^{(2)} = e^{-iz} (2/\pi z)^{1/2} Z_n(1/z)$. When the disk is absent the condition at the infinity demands the relation $b_n = i(-1)^{n+1}a_n$. In the presence of a disk the part of the wave energy can be reflected from a disk and we

have the mixture of both types of waves. In general case the coefficients $a_n^{B,E}, b_n^{B,E}$ are the arbitrary complex numbers and can be determined from the boundary conditions (5,6). The boundary condition on the stellar surface $r = R_s$ ($z = z_0 = R_s \omega_s / c$) are

$$B_r(z = z_0) = B_r(\text{inside}); E_\theta(z = z_0) = E_\theta(\text{inside});$$

$$E_\phi(z = z_0) = E_\phi(\text{inside}).$$

Fields inside are

$$\mathbf{B} = \frac{\mu}{r^3} \sin \chi e^{i\phi - i\omega_s t} \begin{cases} 2P_1^1(\cos \vartheta), & \mathbf{e}_r \\ -\frac{\partial P_1^1}{\partial \vartheta}(1 - \frac{1}{2}z^2), & \mathbf{e}_\theta \\ -i\frac{P_1^1}{\sin \vartheta}(1 - \frac{1}{2}z^2), & \mathbf{e}_\phi, \end{cases} \quad (9)$$

$$\mathbf{E} = \frac{\mu z}{r^3} \sin \chi e^{i\phi - i\omega_s t} \begin{cases} -\frac{1}{3}P_2^1(1 - \frac{1}{2}z^2), & \mathbf{e}_r \\ -\frac{P_1^1}{\sin \vartheta} + \frac{1}{3}\frac{\partial P_2^1}{\partial \vartheta}, & \mathbf{e}_\theta \\ -i\frac{\partial P_1^1}{\partial \vartheta} + i\frac{1}{3}\frac{P_2^1}{\sin \vartheta}, & \mathbf{e}_\phi. \end{cases}$$

Here μ is the magnetic moment of a star. We consider here only the varying component of the magnetic dipole, proportional to $\sin \chi$, which produces an electromagnetic radiation. From (5,6) and (9) it follows

$$(a_1^B J_{3/2}(z) + b_1^B J_{-3/2}(z))|_{z=z_0} = 2\frac{\mu}{R_s^3} z_0^{3/2} \sin \chi;$$

$$(a_n^B J_{n+1/2}(z) + b_n^B J_{-n-1/2}(z))|_{z=z_0} = 0, \quad n \neq 1; \quad (10)$$

$$\left(\frac{\partial}{\partial z} z^{1/2} (a_2^E J_{5/2} + b_2^E J_{-5/2}) \right) \Big|_{z=z_0} = 2\frac{\mu}{R_s^3} z_0^2 \sin \chi;$$

$$\left(\frac{\partial}{\partial z} z^{1/2} (a_n^E J_{n+1/2} + b_n^E J_{-n-1/2}) \right) \Big|_{z=z_0} = 0, \quad n \neq 2.$$

Introducing the notations

$$\varrho_n^B = \frac{J_{n+1/2}(z)}{J_{-n-1/2}(z)} \Big|_{z=z_0}, \quad \varrho_n^E = \frac{\frac{\partial}{\partial z} (z^{1/2} J_{n+1/2})}{\frac{\partial}{\partial z} (z^{1/2} J_{-n-1/2})} \Big|_{z=z_0} \quad (11)$$

and the dimensionless quantities a_n, b_n , dividing them on the value $2\mu \sin \chi / R_s^3$, we rewrite (10) in the form

$$b_1^B = -a_1^B \varrho_1^B + s_1, \quad b_n^B = -a_n^B \varrho_n^B, \quad n \neq 1, \quad (12)$$

$$b_2^E = -a_2^E \varrho_2^E + s_2, \quad b_n^E = -a_n^E \varrho_n^E, \quad n \neq 2.$$

Here the values s_1 and s_2 are

$$s_1 = \frac{z^{3/2}}{J_{-3/2}(z)} \Big|_{z=z_0}, \quad s_2 = \frac{z^2}{\frac{\partial}{\partial z} (z^{1/2} J_{-5/2})} \Big|_{z=z_0}. \quad (13)$$

Let us note that because $z_0 \ll 1$ ($R_s \ll c/\omega_s$) the coefficients $\varrho_n^B, \varrho_n^E, s_1, s_2$ are small values. In the lowest order of z_0 they are

$$\varrho_n^B \approx \frac{\Gamma(1/2 - n)}{\Gamma(3/2 + n)} \left(\frac{z_0}{2} \right)^{2n+1};$$

$$\varrho_n^E \approx -\frac{(n+1)\Gamma(1/2 - n)}{n\Gamma(3/2 + n)} \left(\frac{z_0}{2} \right)^{2n+1};$$

$$s_1 \approx 2^{-3/2} \Gamma(-1/2) z_0^3 = -2^{-1/2} \pi^{1/2} z_0^3;$$

$$s_2 \approx -2^{-7/2} \Gamma(-3/2) z_0^5 = -\frac{2^{-3/2}}{3} \pi^{1/2} z_0^5.$$

Here $\Gamma(x)$ is the Gamma function. In the case of a disk Eq. (12) are the connections between the coefficients b_n and a_n . When a disk is absent connections are conditions $b_n = i(-1)^{n+1} a_n$, that is the condition for the radiation toward the infinity when there are no reflected waves. Then Eq. (12) shows that nonzero coefficients are a_1^B and a_2^E only

$$a_1^B = \frac{(-i + \varrho_1^B) s_1}{(1 + \varrho_1^{B^2})} \approx -i s_1 = i \left(\frac{\pi}{2} \right)^{1/2} z_0^3,$$

$$a_2^E = \frac{(i + \varrho_2^E) s_2}{(1 + \varrho_2^{E^2})} \approx i s_2 = -\frac{i}{3} \left(\frac{\pi}{8} \right)^{1/2} z_0^5.$$

It means that a rotating magnetized star without a disk radiates the magnetodipole radiation with the amplitude $a_1^B \simeq z_0^3$ and also the electroquadrupole radiation with the amplitude $a_2^E \ll a_1^B$. The disk changes this picture, and a star radiates also many multipoles $n \neq 1, 2$.

To find the coefficients $a_n^{B,E}$ we need use the boundary conditions on a disk (5,6). The boundary condition (5) means

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} \left[\frac{P_n^1(\vartheta_0)}{\sin \vartheta_0} z^{-\frac{1}{2}} \frac{\partial}{\partial z} z^{\frac{1}{2}} (a_n^E J_{n+\frac{1}{2}} + b_n^E J_{-n-\frac{1}{2}}) - \frac{\partial P_n^1(\vartheta_0)}{\partial \vartheta_0} (a_n^B J_{n+\frac{1}{2}} + b_n^B J_{-n-\frac{1}{2}}) \right] = 0; \quad z \geq z_d, \quad \vartheta_0 = \frac{\pi}{2} \pm \delta. \quad (14)$$

We use the recurrent relations for the Bessel functions

$$\frac{\partial}{\partial z} z^{\frac{1}{2}} J_{n+\frac{1}{2}} = z^{\frac{1}{2}} \left[\frac{n+1}{2n+1} J_{n-\frac{1}{2}} - \frac{n}{2n+1} J_{n+\frac{3}{2}} \right];$$

$$\frac{\partial}{\partial z} z^{\frac{1}{2}} J_{-n-\frac{1}{2}} = z^{\frac{1}{2}} \left[\frac{n}{2n+1} J_{-n-\frac{3}{2}} - \frac{n+1}{2n+1} J_{-n+\frac{1}{2}} \right].$$

Then multiplying Eq. (14) on the Gegenbauer polynomial $A_{k,1/2}(z)$ and integrating it over the contour passing through the infinity as shown on the figure 2a, we obtain the matrix equation connected the coefficients a_n^B and a_n^E . The Gegenbauer polynomial $A_{k,1/2}$ is the

function conjugated to the Bessel functions $J_{n+1/2}(z)$,

$$A_{k, \frac{1}{2}}(z) = 2^{-\frac{1}{2}}(k + \frac{1}{2}) \sum_{m=0}^{\leq k/2} \frac{\Gamma(\frac{1}{2} + k - m)}{m!} \left(\frac{z}{2}\right)^{2m-k-1},$$

and possesses the property [5]

$$\frac{1}{2i\pi} \int_C J_{n+1/2}(z) A_{k,1/2}(z) z^{-1/2} dz = \delta_{k,n}.$$

The result is

$$\begin{aligned} & \frac{1}{n(n+1)} \frac{\partial P_n^1}{\partial \vartheta_0} (\varrho_n^B C_{k,n} - \delta_{k,n}) a_n^B + \frac{1}{(2n+1)} \frac{P_n^1}{\sin \vartheta_0} \\ & \left[\left(\frac{\delta_{k+1,n}}{n} - \frac{\delta_{n,k-1}}{n+1} \right) + \varrho_n^E \left(\frac{C_{k,n-1}}{n} - \frac{C_{k,n+1}}{n+1} \right) \right] a_n^E = \\ & \frac{1}{2} \frac{\partial P_1^1}{\partial \vartheta} s_1 C_{k,1} + \frac{1}{10} \frac{P_2^1}{\sin \vartheta} s_2 (C_{k,1} - \frac{2}{3} C_{k,3}). \end{aligned} \quad (15)$$

Here coefficients $C_{k,n}$ are

$$C_{k,n} = \frac{1}{2\pi i} \int_C A_{k,1/2} J_{-n-1/2} z^{-1/2} dz =$$

$$(-1)^{l+1} (k + \frac{1}{2}) \sum_{m=0}^{\leq k/2} \frac{(-1)^m \Gamma(\frac{1}{2} + k + m)}{m! (l+1-m)! \Gamma(\frac{1}{2} + k - m - l)}.$$

The coefficients $C_{k,n}$ are not equal to zero if $n+k=2l+1$, l is the integer number.

If we introduce two matrices $\Lambda_{k,n}^{1B}$, $\Lambda_{k,n}^{1E}$ and the vector σ_k^1 ,

$$\begin{aligned} \Lambda_{k,n}^{1B} &= \frac{\partial P_n^1}{\partial \vartheta_0} \frac{1}{n(n+1)} (\varrho_n^B C_{k,n} - \delta_{k,n}); \\ \Lambda_{k,n}^{1E} &= \frac{P_n^1}{\sin \vartheta_0} \frac{1}{(2n+1)} \left[\frac{\delta_{k+1,n}}{n} - \frac{\delta_{k-1,n}}{n+1} + \right. \\ & \left. \varrho_n^E \left(\frac{C_{k,n}}{n} - \frac{C_{k,n+1}}{n+1} \right) \right]; \end{aligned} \quad (16)$$

$$\sigma_k^1 = \frac{1}{2} \frac{\partial P_1^1}{\partial \vartheta_0} s_1 C_{k,1} + \frac{1}{10} \frac{P_2^1}{\sin \vartheta_0} s_2 (C_{k,1} - \frac{2}{3} C_{k,3}),$$

then we obtain

$$\Lambda_{k,n}^{1B} a_n^B + \Lambda_{k,n}^{1E} a_n^E = \sigma_k^1. \quad (17)$$

The second matrix equation connecting a_n^E and a_n^B is followed from the condition (6). Doing the same procedure (multiplying on the Gegenbauer polinomial and integrating in the complex plane z), but taking into account that the Keplerian velocity, $v_k \propto z^{-1/2}$, is the two fold function, we have to introduce the cut in the complex plane from the point $z = z_d$ to the infinity. The resulting contour C has the shape as shown on the

figure 2b.

$$\begin{aligned} & \frac{i\kappa}{\pi} \frac{\partial P_n^1}{\partial \vartheta_0} \frac{\sin \vartheta_0}{2n+1} \left[(d_{k,n-1}^+ + d_{k,n-1}^- \varrho_n^B) \frac{1}{n} - (d_{k,n+1}^+ + \right. \\ & \left. d_{k,n+1}^- \varrho_n^B) \frac{1}{n+1} \right] a_n^B + P_n^1 [\delta_{k,n} - C_{k,n} \varrho_n^E + \\ & \frac{i\kappa}{\pi} \frac{1}{n(n+1)} (d_{k,n}^+ - d_{k,n}^- \varrho_n^E)] a_n^E = \frac{i\kappa}{\pi} \left[\frac{d_{k,0}^+}{3} s_1 \frac{\partial P_1^1}{\partial \vartheta_0} \sin \vartheta_0 \right. \\ & \left. - \frac{d_{k,2}^-}{6} s_2 (P_2^1 + \frac{\partial P_1^1}{\partial \vartheta_0} s_1 \sin \vartheta_0) - P_2^1 s_2 C_{k,2} \right]. \end{aligned} \quad (18)$$

Here

$$d_{k,n}^+ = \int_{z_d}^{\infty} A_{k,1/2}(z) J_{n+1/2}(z) dz, \quad z = z + i0;$$

$$d_{k,n}^- = \int_{z_d}^{\infty} A_{k,1/2}(z) J_{-n-1/2}(z) dz, \quad z = z + i0,$$

$\kappa = (r_g \omega_s / 2c)^{1/2} < 1$ is the Keplerian parameter, r_g is the gravitational radius of a star, $r_g = 2GM_s/c^2$,

$$\kappa = 5.5 \cdot 10^{-3} \left(\frac{M_s}{M_\odot} \right)^{1/2} \left(\frac{P_s}{1s} \right)^{-1/2}.$$

P_s is the period of a star rotation, $P_s = 2\pi/\omega_s$. Again, introducing the matrices $\Lambda_{k,n}^{2B}$, $\Lambda_{k,n}^{2E}$ and the vector σ_k^2 , we rewrite Eq. (18) in the form

$$\Lambda_{k,n}^{2B} a_n^B + \Lambda_{k,n}^{2E} a_n^E = \sigma_k^2; \quad (19)$$

$$\Lambda_{k,n}^{2B} = \frac{i\kappa}{\pi} \frac{\partial P_n^1}{\partial \vartheta_0} \frac{\sin \vartheta_0}{2n+1} \left[(d_{k,n-1}^+ + d_{k,n-1}^- \varrho_n^B) \frac{1}{n} - \right. \\ \left. (d_{k,n+1}^+ + d_{k,n+1}^- \varrho_n^B) \frac{1}{n+1} \right];$$

$$\Lambda_{k,n}^{2E} = P_n^1 \left[\delta_{k,n} - C_{k,n} \varrho_n^E + \frac{i\kappa}{\pi} \frac{1}{n(n+1)} (d_{k,n}^+ - d_{k,n}^- \varrho_n^E) \right]; \quad (20)$$

$$\sigma_k^2 = \frac{i\kappa}{\pi} \left[\frac{d_{k,0}^+}{3} s_1 \frac{\partial P_1^1}{\partial \vartheta_0} \sin \vartheta_0 - \frac{d_{k,2}^-}{6} (s_2 P_2^1 + s_1 \frac{\partial P_1^1}{\partial \vartheta_0} \sin \vartheta_0) \right] - s_2 P_2^1 C_{k,2}.$$

Combining Eq. (17) and Eq. (19) we find that the coefficients a_n^B and a_n^E are the solution of the general matrix equation

$$\Lambda_{k,n} \vec{a} = \vec{\sigma}, \quad (21)$$

where $\vec{a}$ and $\vec{\sigma}$ are the vectors

$$\vec{a} = \begin{pmatrix} a_n^B \\ a_n^E \end{pmatrix}, \quad \vec{\sigma} = \begin{pmatrix} \sigma_k^1 \\ \sigma_k^2 \end{pmatrix}.$$

The matrix $\Lambda_{k,n}$ is

$$\Lambda_{k,n} = \begin{pmatrix} \Lambda_{k,n}^{1B} & \Lambda_{k,n}^{1E} \\ \Lambda_{k,n}^{2B} & \Lambda_{k,n}^{2E} \end{pmatrix}.$$

4 Torque acting on a star

Our purpose is to determine the change of the torque acting on a star under the influence of a rotating disk,

$$\mathbf{K} = \frac{1}{c} \int \mathbf{r} \times \mathbf{J}_s \times \mathbf{B} ds, \quad (22)$$

where $\mathbf{J}_s$ is the surface current on the star surface

$$\mathbf{J}_s = \frac{c}{4\pi r} \times \{\mathbf{B}\}, \quad \{\mathbf{B}\} \equiv \mathbf{B}|_{r=R_s+0} - \mathbf{B}|_{r=R_s-0}. \quad (23)$$

$\{\mathbf{B}\}$ is the discontinuous of the tangential magnetic field. The torque is

$$\mathbf{K} = \frac{R_s}{c} \int \mathbf{J}_s B_{sr} ds, \quad (24)$$

where B_{sr} is the radial magnetic field on the stellar surface

$$B_{sr} = \frac{2\mu}{R_s^3} \sin \chi P_1^1 e^{i\phi - i\omega t}.$$

Integrating in Eq. (24) over the surface we find the component of the torque K_z along the axis of the rotation. It turns out to be proportional to two coefficients of the expansion of the vacuum field over multipoles (7,8)

$$K_z = -\frac{2}{3} \frac{\mu^2 \sin^2 \chi}{R_s^3} \text{Im}(a_1^B) \left[\frac{1}{z} \frac{\partial}{\partial z} (z^{1/2} J_{3/2}) - \frac{1}{z} \frac{\partial}{\partial z} (z^{1/2} J_{-3/2}) \frac{J_{3/2}}{J_{-3/2}} \right]_{z=z_0} + \frac{2}{5} \frac{\mu^2 \sin^2 \chi}{R_s^3} \text{Im}(a_2^E) \left[z^{-1/2} J_{5/2} - z^{-1/2} J_{-5/2} \frac{\frac{\partial}{\partial z} (z^{1/2} J_{5/2})}{\frac{\partial}{\partial z} (z^{1/2} J_{-5/2})} \right]_{z=z_0}. \quad (25)$$

That means that only two types of radiation, the magnetodipole and the electroquadrupole, retard the star rotation. The contribution of the magnetodipole component is much larger than the electroquadrupole one because $z_0 \ll 1$. In the absence of a disk the torque is

$$K_z = K_z^{MD} = -\frac{2}{3} \frac{\mu^2 \sin^2 \chi}{R_s^3} z_0^3 = -\frac{2}{3} \frac{\mu^2 \omega^3 \sin^2 \chi}{c^3}, \quad (26)$$

$$|K_z^{MD}| = 1.5 \cdot 10^{30} B_{12}^2 \left(\frac{R}{10 \text{ km}} \right)^6 \left(\frac{P_s}{1 \text{ s}} \right)^{-3} \sin^2 \chi \text{ erg}.$$

An conducting disk distorts the radiation. There appears another multipole components of the radiation. The amplitudes of them are connected by the relation (21). The coefficients $\Lambda_{k,n}$ contain also the imaginary parts which are proportional to the Keplerian parameter κ . The disk also changes the conditions at the infinity - a disk reflects the wave energy. As a result a star can as lost the angular momentum as obtain it from a disk. The effect depends on the disk parameter $z_d = r_d \omega_s / c$. The qualitative estimation demands

the solution of the matrix equation (21) numerically. The result of calculations for the usual neutron star parameters $R = 10 \text{ km}$, $M_s = 1 M_\odot$, $\omega_s = 1 \text{ s}^{-1}$ and $z_d = 1$, $\delta = 10^\circ$ is presented on the table 1.

n	1	2	3	4	5	6
$\text{Im} a_n^B$	0.25	0.15	0.05	0.013	0.004	0.0027
$\text{Re} a_n^B$	0.21	0.18	0.05	0.013	0.008	0.003
$\text{Im} a_n^E$	0.25	0.20	0.075	0.027	0.007	0.0027
$\text{Re} a_n^E$	0.25	0.28	0.065	0.027	0.008	0.0027

Table 1. Amplitudes of n-th magnetic multipoles a_n^B and n-th electric multipoles a_n^E for the first sixth waves (7,8). They are normalized by their values in the vacuum: a_n^B by $\text{Im} a_1^B = (\pi/2)^{1/2} z_0^3$ and a_n^E by $\text{Im} a_2^E = -(\pi/72)^{1/2} z_0^5$.

We see that though the main component is a_1^B , there appear compatible amplitudes a_2^B, a_3^B, a_4^B . Among electric multipoles the electro-dipole amplitude a_1^E becomes general. For the larger $n > 4$ the amplitudes of multipoles fall down. It permits us to restrict ourself by the finite dimension of vectors $\vec{d}$, $\vec{\sigma}$ in equation (21), $k = 12$. The calculation of the torque K_z , acting on a star, is presented on the figure 3. For comparison we draw also the result of calculation for $k = 8$. It is seen that the difference between $k = 8$ and $k = 12$ is not significant and our approximation to use finite numbers of equations in infinite matrix equation (21) is valid. The main result is the change of sign of K_z at $z_d = z^* \approx 1$. For the chosen parameters $z^* = 0.8$. The behavior of the torque K_z over z_d can be well approximated the by the simple formula

$$K_z = K_z^{MD} \left(1 - \frac{z^{*2}}{z_d^2} \right). \quad (27)$$

This expression does not take into account the small maximum of K_z at $z_d \approx 2 - 3$, when a star transmits its angular momentum not only to the electromagnetic radiation, but also to a disk.

At $z_d < z^*$ a disk transmits the angular momentum to a star, it spins up. Let us note that $z^* \approx 1$ is just the region where the electromagnetic radiation forms. A disk changes radiation conditions in the zone where it originates. Such electromagnetic stellar spin up exists independently of the mechanical angular momentum transition, which is proportional to the value of the accretion matter rate $\dot{M}$.

The structure of the magnetic field is shown on the figure 4. Here we present only the magnetic field produced by the varying component of the magnetic dipole, proportional to $\mu \sin \chi$, which generates the electromagnetic radiation.

We see that the magnetic field lines repeal from an ideal disk. It is the result of calculations under the




Fig. 2 The contour of integration in the complex plane z . The contour a) corresponds to the boundary condition (5), and the contour b) corresponds to the boundary condition (6) on the disk surface.




Fig. 3 The torque K_z , acting on a star, in the units of K_z^{MD} versus the inner edge of a disk z_d . $\delta = 10^\circ$, a) $k=12$, b) $k=8$.




Fig. 4 Projection of magnetic field lines of the oblique dipole onto the plane orthogonal to the direction of the magnetic moment μ . Owing to the projective view and the large scale of the order of light cylinder radius c/ω_s , some details on this figure look like peculiarities (cusps, touches), but they are not in reality.

boundary conditions (5,6) on the disk surface. It is not postulated ad hoc as did in [2].

5 Discussion

We showed that presence of a conducting disk in the magnetosphere of a rotating magnetized star changes significantly the electromagnetic radiation of a star if the inner edge of a disk r_d is closer than the radius of the stellar light cylinder c/ω_s . The torque acting on a star K_z changes its sign when $r_d < r^* \simeq c/\omega_s$. Instead of the spinning down of the star rotation, a star begins to spin up, getting the angular momentum from a disk. At the position of the inner disk radius near the light cylinder radius, $r_d = r^*$, the torque K_z becomes zero. It means that a star will not change its angular momentum and the period of its rotation will be constant. It can happen after the period of the gas accretion onto a star when it gains the angular momentum, value of which is proportional to the accretion rate $\dot{M}$. As was shown [6], under the mass accretion onto a magnetized star, the torque K_z has always definite sign, and there is no situation of the stable star rotation. When the accretion stops, $\dot{M} = 0$, a star pushes out a disk from its vicinity, that is, so called, the propeller regime. The inner edge of a disk moves toward the light cylinder distance where a disk prevents the stellar spin down due to the radiation of electromagnetic waves. The period of the stellar rotation will not change when a disk is presented at the distance from a star $r_d \simeq c/\omega_s$. Such period of the star life is observed among X-rays binaries [1].

Acknowledgements. This work was partially supported by Russian Foundation for Basic Research (Grant no. 11-02-01021).

References

- [1] Lipunov V.M., 1993, *Astrophysics of Neutron Stars*. Springer-Verlag, Berlin.
- [2] Aly J.J., 1980, *Astron. Astrophys.* **86**, p. 192-197.
- [3] Bardou A., Heyvaerts J., 1996, *Astron. Astrophys.* **307**, p.1009- 1022.
- [4] Lovelace R.V.E., Romanova M.M., Bisnovatyi-Kogan G.S., 1995, *MNRAS*, **275**, p. 244-254.
- [5] Watson G.N., *A treatise on the theory of Bessel functions*, 1966, Cambridge University Press, p. 283, 524, 525.
- [6] Istomin Ya.N., Haensel P., 2012, *MNRAS*, in press.